

Short communication

## Chromosome numbers of South African Umbelliferae (Apiaceae)

J.V. Shner<sup>a</sup>, T.V. Alexeeva<sup>a</sup>, M.G. Pimenov<sup>a</sup>, B.-E. Van Wyk<sup>b,\*</sup>

<sup>a</sup> Botanical Garden, Moscow State University, Moscow 119899, Russia

<sup>b</sup> Department of Botany and Plant Biotechnology, University of Johannesburg, Johannesburg, South Africa

Received 31 December 2009; received in revised form 15 July 2010; accepted 19 October 2010

### Abstract

Chromosome numbers are reported for 12 species from nine genera of South African Umbelliferae, of which seven species and one genus (*Itasina*) are recorded for the first time. A detailed list of all published chromosome counts for southern African species is also presented, together with a review of the literature. The new data obtained are briefly discussed in the context of the taxonomy and relationships of local Umbelliferae. The counts agree with previous reports except that *Annesorhiza* appears to have  $2n=22$ , with or without one additional B-chromosome, and not  $2n=24$  as reported in the literature. The number for *Itasina* ( $2n=24$ ) is of considerable interest and indicates that a detailed chromosome study of the South African genera *Annesorhiza* and *Chamarea* may yield valuable taxonomic information.

© 2010 SAAB. Published by Elsevier B.V. All rights reserved.

**Keywords:** *Anginon*; Apiaceae; Apioideae; Cape Floristic Region; Chromosome counts; *Itasina*

### 1. Introduction

Chromosome numbers are part of a multidisciplinary approach to the characterization of taxa. The South African Umbelliferae include several anomalous and taxonomically isolated genera that are of importance in unravelling the relationships within the family as a whole (Burt, 1991; Magee et al., 2010a,b; Van Wyk and Tilney, 2004), yet chromosome counts are available only for a limited number of species. Among the approximately 205 species of southern African Umbelliferae (Allison and Van Wyk, 1997; Burt, 1991; Liu et al., 2007; Magee et al., 2008a,b, 2009a,b; Schubert and Van Wyk, 1997; Tilney and Van Wyk, 2002; Van Wyk and Tilney, 2003, 2004; Winter et al., 2008), chromosome counts based on local material are available for 25 species only, belonging to 19 genera (Table 1). Most of them are South African endemics or they are at least endemic or subendemic to the African continent. The only exceptions are the more widely distributed *Apium graveolens* L. and *Torilis arvensis* (Huds.) Link. This means that chromosome numbers of only 23 South African

species (indigenous or naturalised) have been determined up to now (Pimenov et al., 2002).

The most important sources of data include two publications by L. Constance et al. (Constance and Chuang, 1982; Constance et al., 1976). Some additional references on the identity of the species that were counted can be found in Burt (1991). The only other counts reported are those of Riley and Hoff (1961) for *Steganotaenia araliacea* Hochst., Cauwet (1971) and Cauwet-Marc (1979) for *Bupleurum mundtii* Cham. & Schtdl. and Winter and Van Wyk (1996) for some South African species of *Heteromorpha* Cham. & Schtdl.

The aim of this paper is to (1) report chromosome numbers for an additional 12 South African species from nine genera of Umbelliferae, and (2) to review all available data on southern African species and the associated literature.

### 2. Materials and methods

Plant material (flower buds and fruits) was collected during two short excursions before and after the 4th International Apiales Symposium in Pretoria, 3 to 19 January 2003 (Table 2). Author citations for the scientific names given in Tables 1 and 2 are not repeated from here on. Fixations were made by M.G. Pimenov and voucher specimens were collected and identified

\* Corresponding author. Tel.: +27 115592412; fax: +27 115592411.  
E-mail address: [bevanwyk@uj.ac.za](mailto:bevanwyk@uj.ac.za) (B.-E. Van Wyk).

by M.G. Pimenov, B.-E. Van Wyk, E.V. Kljuykov and T.A. Ostroumova; they are kept in MW.

Meiotic chromosomes were examined in pollen mother cells (Fig. 1). Flower buds were fixed in the field in 3:1 96% ethanol–glacial acetic acid mixture and then stained with 1% acetocarmine for 3 to 4 days at about 5 °C. After staining, the anthers were squashed and mounted. Somatic chromosomes were examined in meristem cells of root tips from germinating seeds (Fig. 2). After a pretreatment in 0.01% aqueous colchicine solution for 3 h at room temperature, the root tips were fixed in 3:1 96% ethanol–glacial acetic acid mixture and then stained with 1% acetocarmine at 100 °C for 3 to 5 min. Both the anthers and the root tips were squashed on a glass microscope slide in chloral hydrate (Hoyer's solution).

### 3. Results and discussion

Chromosome numbers were determined for 12 species from nine genera (Table 2). These counts include a first record for the genus *Itasina* Raf. and also first counts for seven other species (*Afrosciadium caffrum*, *A. magalismontanum*, *Anginon difforme*, *A. paniculatum*, *A. swellendamense*, *Lichtensteinia lacera* and *Notobubon capense*). In addition, a first correct count for *Annesorhiza macrocarpa* is presented.

The chromosome number of *Itasina filifolia* ( $n=12$ ) is rather unusual in Apioideae. It is known for only 24 species of the subfamily (representing 1.6% of species investigated) and was found in the traditional tribes Scandiceae (two species), Caucalideae (four species), Smyrnieae (one species), Apieae (14 species), Peucedaneae (one species) and Tordylieae (two species). *Itasina* traditionally belongs to the tribe Apieae but molecular systematic evidence recently showed that it is related to the genera *Annesorhiza* Cham. & Schldl., *Astydamia* DC., *Chamarea* Eckl. & Zeyh., *Ezosciadium* B.L.Burt and *Molopospermum* W.J.D. Koch (Magee et al., 2010b) and that these genera should be recognized as a new tribe, the Annesorhizeae Magee et al. (Magee et al., 2010b). These genera (especially *Ezosciadium*) would be worth sampling in further cytological studies. Counts of  $n=12$  (or  $2n=24$ ) occur sporadically in species of unrelated genera, including *Ammoselinum* Torr. & A. Gray, *Apium* L., *Bunium* L., *Bupleurum* L., *Carum* L., *Cnidium* Cusson ex Jussieu, *Pimpinella* L., *Spuriopimpinella* (H. Boisseu) Kitag. and *Tilingia* Regel (Pimenov et al., 2002). The count of  $n=12$  in Umbelliferae deserves special attention as it is common in Araliaceae, the reputedly related family. Moreover,  $x=12$  or  $x=6$  is regarded as the most plausible basic and ancestral chromosome number in Apiales (Sharma and Chatterji, 1964; Yi et al., 2004). In connection with this putative close affinity,  $n=12$  in the woody African species *S. araliacea*

Table 1  
Previously determined chromosome numbers for 19 genera and 25 species of South African Umbelliferae.

Genera and species	Chromosome number	References
<i>Afrologisticum wilmsianum</i> (H. Wolff) P.J.D.Winter [as <i>Peucedanum wilmsianum</i> H.Wolff]	$n=11$	Constance and Chuang (1982)
<i>Afrosciadium platycarpum</i> (Sond.) P.J.D.Winter [as <i>Peucedanum platycarpum</i> E.Mey ex Sond.]	$n=22$	Constance and Chuang (1982)
	$n=11$	Burt (1991) [Eastern Cape specimen]
<i>Agrocharis melanantha</i> Hochst.	$n=22$	Constance and Chuang (1982); Burt (1991, p. 152; Sani Pass specimen)
<i>Alepidea natalensis</i> J.M.Wood & Evans	$n=8$	Constance and Chuang (1982)
<i>Anginon rugosum</i> (Thunb.) Raf. [= <i>A. uitenhagense</i> (Eckl. & Zeyh.) B.L.Burt], as	$n=11$	Burt (1991)
<i>Annesorhiza macrocarpa</i> Eckl. & Zeyh.	$n=12$	Burt (1991)
<i>Apium graveolens</i> L.	$n=11$	Constance et al. (1976); Constance and Chuang (1982)
<i>Arctopus echinatus</i> L.	$n=9$	Constance et al. (1976)
<i>Bupleurum mundtii</i> Cham. & Schldl.	$2n=16$	Cauwet (1971); Cauwet-Marc (1979)
<i>Conium chaerophylloides</i> (Thunb.) Sond.	$n=11$	Constance and Chuang (1982)
<i>C. fontaneum</i> Hilliard & B.L.Burt	$n=11$	Constance and Chuang (1982)
<i>Cynorhiza typica</i> Eckl. & Zeyh. [as <i>Peucedanum olifantianum</i> (Koso-Pol.) M.Hiroe]	$n=11$	Constance and Chuang (1982)
<i>Dasispermum capense</i> (Lam.) Magee & B.-E.Van Wyk [as <i>Stoibrax capense</i> (Lam.) B.L.Burt]	$n=9$	Constance et al. (1976)
<i>D. humile</i> (Meisn.) Magee & B.-E.Van Wyk [as <i>Sonderina humilis</i> (Meisn.) H.Wolff]	$n=8$	Burt (1991)
<i>D. suffruticosum</i> (Berg.) B.L.Burt	$n=9$	Constance et al. (1976); Constance and Chuang (1982)
<i>Glia prolifera</i> (Burm.f.) B.L.Burt	$n=11$	Constance et al. (1976)
<i>Hermas villosa</i> (L.) Thunb.	$n=7$	Constance et al. (1976)
<i>Heteromorpha arborescens</i> (Spreng.) Cham. & Schldl.	$n=11$	Constance and Chuang (1982)
<i>H. arborescens</i> var. <i>frutescens</i> P.J.D.Winter	$2n=22$	Winter and Van Wyk (1996)
<i>H. involucrata</i> Conrath	$2n=22$	Winter and Van Wyk (1996)
<i>H. pubescens</i> Burt Davy	$2n=22$	Winter and Van Wyk (1996)
<i>Lichtensteinia interrupta</i> (Thunb.) Sond. [as <i>L. kolbeana</i> Bolus]	$n=11$	Constance and Chuang (1982)
<i>Notobubon galbanum</i> (L.) Magee [as <i>Peucedanum galbanum</i> (L.) Benth. & Hook.f.]	$n=11$	Constance and Chuang (1982)
<i>N. laevigatum</i> (Aiton) Magee [as <i>Peucedanum capense</i> (Thunb.) Sond.]	$n=11$	
<i>Steganotaenia araliacea</i> Hochst.	$2n=22$	Riley and Hoff (1961); Constance and Chuang (1982)
<i>Torilis arvensis</i> (Huds.) Link	$n=6$	Constance et al. (1976)

Table 2

Newly determined chromosome numbers for nine genera and 12 species of South African Umbelliferae (specimens are housed in MW).

Species	Origin and voucher specimen	n	2n
<i>Afroscidium caffrum</i> (Meisn.) P.J.D.Winter [= <i>Peucedanum caffrum</i> (Meisn.) Philips]	Limpopo Province, slopes of Wolkberg, Haenertsburg, 23°56'S, 29°57'E. 05.01.2003, Pimenov et al. 17		22
<i>A. magalimontanum</i> (Sond.) P.J.D.Winter [= <i>Peucedanum magalimontanum</i> Sond.]	Gauteng Province, 20 km E of Pretoria, near Bronkhorstspuit, 25°46'S, 28°32'E. 03.01.2003, Pimenov et al. 1		22
<i>Anginon difforme</i> (L.) B.L.Burt	Western Cape Province, Kirstenbosch, 34°03'S, 18°23'E. 14.01.2003, Pimenov et al. 73	11	
<i>A. paniculatum</i> (Thunb.) B.L.Burt	Western Cape Province, near Citrusdal, 32°25'S, 18°57'E. 15.01.2003, Pimenov et al. 82	11	
<i>A. swellendamense</i> (Eckl. & Zeyh.) B.L.Burt	Western Cape Province, Worcester, 33°12'S, 19°23'E. 15.01.2003, Pimenov et al. 94	11	
<i>Annesorhiza macrocarpa</i> Eckl. & Zeyh.	Western Cape Province, Noordhoek, 34°05'S, 18°21'E. 13.01.2003, Pimenov et al. 59	11, 11+1B	
<i>Cyclosporum leptophyllum</i> (Pers.) Sprague ex Britton & P.Wilson	Limpopo Province, slopes of Wolkberg, Haenertsburg, 23°56'S, 29°57'E. 05.01.2003, Pimenov et al. 13		14
<i>Dasispermum suffruticosum</i> (Berg.) B.L.Burt	Western Cape Province, Betty's Bay, Stony Point, 34°21'S, 18°55'E. 12.01.2003, Pimenov et al. 47	9	18
<i>Hermas villosa</i> (L.) Thunb.	Western Cape Province, Table Mountain, 33°57'S, 18°24'E. 13.01.2003, Pimenov et al. 67	7	
<i>Itasina filifolia</i> (Thunb.) Raf.	Western Cape Province, Cape Peninsula, 34°21'S, 18°22'E. 3.01.2003, Pimenov et al. 61	12	
<i>Lichtensteinia lacera</i> Cham. & Schtdl.	Western Cape Province, near Silver Mine, 34°05'S, 18°25'E. 13.01.2003, Pimenov et al. 49	11	
<i>Notobubon capense</i> (Eckl. & Zeyh.) A.R.Magee [= <i>Peucedanum multiradiatum</i> Drude]	Western Cape Province, Betty's Bay, 34°21'S, 18°55'E. 12.01.2003, Pimenov et al. 41		22

(Constance and Chuang, 1982) could be of considerable interest. *Steganotaenia* and *Polemanniopsis* B.L. Burt (both woody shrubs or trees) have recently been recognized as a new tribe, the Steganotaeniae C.I. Calviño & S.R. Downie (Calviño and Downie, 2007). The count of  $n=12$  in the herbaceous *Itasina* however, seems to be of independent origin, as there is no other evidence of a direct connection with *Steganotaenia* Hochst. or with Araliaceae. It is interesting to note that *Itasina* and *Steganotaenia* both form part of the early diverging and predominantly African lineages of the subfamily Apioideae (Magee et al., 2010b).

Our determination does not confirm the reported chromosome number of  $n=12$  for *A. macrocarpa*, published by Burt (1991) with a reference to Constance and Chuang, cited as “ $n=12$  (C & C ined. based on *Batten 590* from E. Cape, E. London). Our data showed that the species has  $n=11$ , but sometimes its karyotype contains one additional B-chromosome. It is therefore likely that earlier researchers incorrectly counted a B-chromosome as one of the basic set. Since *Annesorhiza*, *Chamarea* and *Itasina* are now considered to be closely related (Magee et al., 2010b; Vessio, 2001), a wider survey of these genera seems desirable and will yield valuable information.

There is no straightforward evidence for the chromosome number of any species of the woody genus *Anginon* Raf. in the karyological literature. However, Burt (1991) showed  $n=11$  for *A. uitenhagense* on the basis of a re-identification of the *Bayliss 8877* collection studied by Constance and Chuang (1982) and reported as “*Peucedanum zeyheri*”. Allison and Van Wyk (1997) reduced *A. uitenhagense* to the synonymy of *A. rugosum* but did not cite *Bayliss 8877* among the material examined. We determined the same count of  $n=11$  for three correctly identified species of *Anginon* (see Table 2). This means that all counts available for the genera of the tribe Heteromorphae M.F. Watson & S.R. Downie (*Anginon*, *Glia* Sond. and *Heteromorpha*) are  $n=11$  or  $2n=22$ .

The chromosome number of *Cyclosporum leptophyllum* [= *Apium leptophyllum* (Pers.) F.Muell.], a widely distributed

weed of warm and adjacent temperate regions of the world, has been determined at least 12 times. The determinations for plants from South and North America, Oceania, Europe and China invariably showed a count of  $n=7$  (or  $2n=14$ ). This was confirmed by the present determination, the first one for African material.

For *Dasispermum suffruticosum* and *Hermas villosa*, our counts confirm previous ones ( $n=9$  and  $n=7$  respectively). In the first case, the chromosome number distinguishes *Dasispermum* Neck. ex Raf. from the genera *Conium* L., *Athamanta* L., *Cicuta* L., *Cnidium*, *Selinum* L., *Sium* L. and others, to which the species has been referred to at various times during its long taxonomic history. Recently, Magee et al. (2009b) have shown, on the basis of morphological and molecular evidence, that *Dasispermum* is congeneric with another South African genus *Sonderina* H. Wolff, and that *Stoibrax capensis* is also part of this genus. The known counts of  $n=8$  for *Dasispermum humile* (= *Sonderina humilis*) and  $n=9$  for *D. capense* (= *S. capensis*) (Table 1) indicate that the more broadly circumscribed *Dasispermum* has  $n=8$  or  $n=9$  and that there is a reduction in the chromosome number in this group. More counts will be valuable and interesting, as the genus now comprises seven species (Magee et al., 2009b, 2010a).

*Hermas* L. is an anomalous genus with affinities in the new subfamily Azorelloideae, where it was moved from its traditional position in the subfamily Hydrocotyloideae. In general, the subfamily Hydrocotyloideae in its traditional circumscription is not monophyletic (Nicolas and Plunkett, 2009). Within the subfamily Azorelloideae, the count of  $n=7$  found in *Hermas*, is unique. Traditionally *Hermas* had been included in the tribe of Mulineae, but modern molecular analyses (Nicolas and Plunkett, 2009; Plunkett, 2001) showed an isolated position for the tribe on the border between Umbelliferae and Araliaceae. Morphological analyses by Liu (2004) and Liu et al. (2009), as well as molecular systematic evidence (Plunkett et al., 2004) indicated that the group should be recognized as a separate subfamily, the Azorelloideae.




Fig. 1. Microphotographs of meiotic chromosomes of some South African Umbelliferae. (a) *Anginon difforme* (n=11); (b) *A. paniculatum* (n=11); (c) *A. swellendamense* (n=11); (d) *Annesorhiza macrocarpa* (n=11); (e) *A. macrocarpa* with B-chromosomes (n=11+1B) [a second set of chromosomes that did not separate is visible to the right, with their corresponding B-chromosome]; (f) *Dasispermum suffruticosum* (n=9); (g) *Hermas villosa* (n=7); (h) *Itasina filifolia* (n=12); and (i) *Lichtensteinia lacera* (n=11). Voucher specimens are listed in Table 2. Scale bar=10  $\mu$ m.

The number of  $n=11$  for *L. lacera* is the second determination for the genus. It corresponds to Constance and Chuang's (1982) count for the eastern Cape form of *L. interrupta*, previously known as *L. kolbeana*. This number is widely distributed in Apioideae, but very rare in Saniculoideae (only the critical South American tetraploid genus *Oligocladus* Chodat & Wilczek has  $n=11$ ). This confirmation is important because it strongly supports the idea that *Lichtensteinia* is more closely related to the subfamily Apioideae than to the Saniculoideae, despite sharing several morphological characters with the latter. Molecular evidence indicates that *Lichtensteinia* is sister to the cytologically unknown genus *Marlothiella* H. Wolff and that both are sister to all other Apioideae (Calviño et al., 2006, 2008; Magee et al., 2010b).

Finally, according to our data, three South African species that were previously included in the broad concept of *Peucedanum* L. have  $2n=22$ . The two herbaceous species from the northern part of South Africa are now included in the newly described genus *Afroscidium* P.J.D.Winter, while the shrubby species from the Cape are now placed in the genus *Notobubon* B-E.Van Wyk (Magee et al., 2009a; Winter et al., 2008). These data are unfortunately not very informative to elucidate relationship among the species, because this count is the most frequent in *Peucedanum* sensu lato as well as in related and unrelated genera of Apioideae. The need to subdivide African *Peucedanum* species was also indicated by the studies

of Ostroumova and Pimenov (1997), where the considerable diversity in the fruit structure was highlighted for the first time. We were unable to confirm the occurrence of tetraploidy ( $n=22$ ), reported in *Afroscidium platycarpum* from the eastern Cape and also *Agrocharis melanantha* from Sani Pass (Table 1).

#### 4. Conclusions

Chromosome numbers of the Umbelliferae of South Africa are insufficiently known and do not yet allow for global comparisons of the family as a whole. We determined chromosome numbers for 12 Umbelliferae species of the South African flora. The genus *Itasina* and seven other species were counted for the first time while the previously reported chromosome numbers of the remaining species were confirmed or sometimes corrected. This study and the review of available data showed that chromosome information can be of value in critical taxonomic revisions of the interesting and anomalous early diverging lineages of southern African Umbelliferae.

#### Acknowledgements

Our cordial thanks are addressed to all organizers and participants of the 4th International Apiales Symposium and its excursions. The Russian authors acknowledge the RFFI; MGP






Fig. 2. Microphotographs of somatic chromosomes of some South African Umbelliferae. (a) *Cyclospermum leptophyllum* ( $2n=14$ ); (b) *Dasispermum suffruticosum* ( $2n=18$ ); (c) *Afrosciadium caffrum* ( $2n=22$ ); (d) *A. magalimontanum* ( $2n=22$ ); and (e) *Notobubon capense* ( $2n=22$ ). Voucher specimens are listed in Table 2. Scale bar = 10  $\mu\text{m}$ .

and BEVW additionally thank the University of Johannesburg and the National Research Foundation for financial support.

## References

- Allison, I., Van Wyk, B.-E., 1997. A revision of the genus *Anginon* (Apiaceae). *Nordic Journal of Botany* 17, 561–577.
- Burt, B.L., 1991. Umbelliferae of southern Africa: an introduction and annotated checklist. *Edinburgh Journal of Botany* 48, 133–282.
- Calviño, C.I., Downie, S.R., 2007. Circumscription and phylogeny of Apiaceae subfamily Saniculoideae based on chloroplast DNA sequences. *Molecular Phylogenetics and Evolution* 44, 175–191.
- Calviño, C.I., Tilney, P.M., Van Wyk, B.-E., Downie, S.R., 2006. A molecular phylogenetic study of southern African Apiaceae. *American Journal of Botany* 95, 196–214.
- Calviño, C.I., Martínez, S.G., Downie, S.R., 2008. Morphology and biogeography of Apiaceae subfamily Saniculoideae as inferred by phylogenetic analysis of molecular data. *American Journal of Botany* 93, 1832–1833.
- Cauwet, A.-M., 1971. Contribution à l'étude caryosystématique du genre *Bupleurum* L. III. *Bulletin de la Société botanique de France* 114, 55–68.
- Cauwet-Marc, A.-M., 1979. Connaissances caryologiques actuelles sur le genre *Bupleurum* L. (Umbelliferae): nombres chromosomiques et nombres de base. *Bulletin de Musée National d'Histoire Naturelle: 4e série Botanique*, vol. 1, pp. 191–211.
- Constance, L., Chuang, T.-I., 1982. Chromosome numbers of Umbelliferae (Apiaceae) from Africa south of the Sahara. *Botanical Journal of the Linnean Society* 85, 195–208.
- Constance, L., Chuang, T.-I., Bell, S.R., 1976. Chromosome numbers in Umbelliferae. IV. *American Journal of Botany* 63, 608–625.
- Liu, M., 2004. A taxonomic evaluation of fruit structure in the family Apiaceae. PhD Thesis, University of Johannesburg.
- Liu, M., Van Wyk, B.-E., Tilney, P.M., 2007. A taxonomic revision of the genus *Choritaenia* (Apiaceae). *South African Journal of Botany* 73, 184–189.
- Liu, M., Van Wyk, B.-E., Tilney, P.M., Plunkett, G.M., Lowry II, P.P., 2009. Evidence from fruit structure supports in general the circumscription of Apiaceae subfamily Azorelloideae. *Plant Systematics and Evolution* 280, 1–13.
- Magee, A.R., Van Wyk, B.-E., Tilney, P.M., Van der Bank, M., 2008a. A taxonomic revision of the South African endemic genus *Arctopus* (Apiaceae, Saniculoideae). *Annals of the Missouri Botanical Garden* 95, 471–486.
- Magee, A.R., Tilney, P.M., Van Wyk, B.-E., Downie, S.R., 2008b. *Ezoscium*: yet another early diverging South African apioid. *Plant Systematics and Evolution* 276, 167–175.
- Magee, A.R., Tilney, P.M., Van Wyk, B.-E., 2009a. A revision of the woody Cape genus *Notobubon* (Apiaceae). *Systematic Botany* 34, 220–242.
- Magee, A.R., Tilney, P.M., Van Wyk, B.-E., Downie, S.R., 2009b. Generic delimitations and relationships of the Cape genera *Capnophyllum*, *Dasispermum* and *Sonderina*, the North African *Kruberia* and *Stoibrax*, and a new monotypic genus of the subfamily Apioideae (Apiaceae). *Systematic Botany* 34, 580–594.
- Magee, A.R., Van Wyk, B.-E., Tilney, P.M., Downie, S.R., 2010a. A taxonomic revision of the genus *Dasispermum* (Apiaceae, Apioideae). *South African Journal of Botany* 76, 308–323.
- Magee, A.R., Calviño, C.I., Liu, M., Downie, S.R., Tilney, P.M., Van Wyk, B.-E., 2010b. New tribal delimitations for the early diverging lineages of Apiaceae subfamily Apioideae. *Taxon* 59, 567–580.
- Nicolas, A.N., Plunkett, G.M., 2009. The demise of subfamily Hydrocotylodeae (Apiaceae) and the re-alignment of its genera across the entire order Apiales. *Molecular Phylogenetics and Evolution* 53, 134–151.

- Ostroumova, T.V., Pimenov, M.G., 1997. Carpological diversity of African *Peucedanum* s.l. I. The species of southern Africa. Feddes Repertorium 108, 299–318.
- Pimenov, M.G., Vassil'eva, M.G., Daushkevich, J.V., Leonov, M.V., 2002. Karyotaxonomical analysis in the Umbelliferae. Science Publishers, Enfield (USA), p. 468.
- Plunkett, G.M., 2001. Relationship of the order Apiales to subclass Asteridae: re-evaluation of morphological characters based on insights from molecular data. Edinburgh Journal of Botany 58, 183–200.
- Plunkett, G.M., Chandler, G.T., Lowry, P.P., Pinney, S., Sprenkle, T., 2004. Recent advances in understanding Apiales with a revised classification. South African Journal of Botany 70, 371–381.
- Riley, H.P., Hoff, V.J., 1961. Chromosome studies in some South African Dicotyledons. Canadian Journal of Genetics and Cytology 3, 260–271.
- Schubert, M.T.R., Van Wyk, B.-E., 1997. A revision of *Centella* series *Capenses* (Apiaceae). Nordic Journal of Botany 17, 301–314.
- Sharma, A.K., Chatterji, A.K., 1964. Cytological study as an aid in the interpretation of the systematic status of different genera of Araliaceae. Cytologia 29, 1–2.
- Tilney, P.M., Van Wyk, B.-E., 2002. A revision of the genus *Annesorhiza* (Apiaceae). Nordic Journal of Botany 21, 615–649.
- Van Wyk, B.-E., Tilney, P.M., 2003. Apiaceae. In: Germishuizen, G., Meyer, N.L. (Eds.), Plants of southern Africa: an annotated checklist. : Strelitzia, vol. 14. National Botanical Institute, Pretoria, pp. 123–132.
- Van Wyk, B.-E., Tilney, P.M., 2004. Diversity of Apiaceae in Africa. South African Journal of Botany 70, 433–445.
- Vessio, N., 2001. The generic affinities of deciduous species of the genera *Annesorhiza* Cham. & Schlecht., *Chamarea* Eckl. & Zeyh. and *Peucedanum* L. (Apiaceae). MSc dissertation, University of Johannesburg.
- Winter, P.J.D., Van Wyk, B.-E., 1996. A revision of the genus *Heteromorpha* (Apiaceae). Kew Bulletin 51, 225–265.
- Winter, P.J.D., Magee, A.R., Phephu, N., Tilney, P.M., Downie, S.R., Van Wyk, B.-E., 2008. A new generic classification for African peucedanoid species (Apiaceae). Taxon 57, 347–364.
- Yi, T., Lowry II, P.P., Plunkett, G.M., Wen, J., 2004. Chromosomal evolution in Araliaceae and close relatives. Taxon 53, 987–1005.